



GOVERNMENT OF THE NORTHWEST TERRITORIES

TECHNICAL DATA SUMMARY
ARSENIC IN THE YELLOWKNIFE ENVIRONMENT
YELLOWKNIFE, NORTHWEST TERRITORIES

REVISED EDITION

JULY 1977

NWT
EX
EPEP
Tec
1977b




Technical Data Summary

Arsenic in the Yellowknife Environment

Yellowknife, Northwest Territories.

Revised Edition

June, 1977.



Prepared by the Ad Hoc
Standing Committee
on Arsenic

D. Billing, Chairman.
Yellowknife, N.W.T.

Compiled and edited by D.A. Gemmill, P. Eng.

Abstract

This report summarizes all of the available information on the level of arsenic in various segments of the Yellowknife, N.W.T., environment. The data was obtained from several studies conducted by a number of federal government departments (Fisheries and the Environment, Indian Affairs and Northern Development, National Health and Welfare), the Government of the Northwest Territories, and other agencies.

A summary of arsenic levels is presented below :

1. Concentrations in the air averaged $0.1 \text{ ug}/\text{m}^3$ (micrograms per cubic meter) in the City and $0.3 \text{ ug}/\text{m}^3$ at Giant Mine.
2. Ambient arsenic levels in suspended particulates in the air over the City were, with few exceptions, always $<$ (less than) $0.1 \text{ ug}/\text{m}^3$.
3. Arsenic concentrations in the Yellowknife municipal water supply were, with very few exceptions, always $< 0.01 \text{ mg}/\text{l}$ (milligrams per liter), the desirable National Standard.
4. Arsenic concentrations in Back Bay, a part of which is the potable mill water source for Giant Mine, ranged from $< 0.01 \text{ mg}/\text{l}$ to $>$ (greater than) $0.7 \text{ mg}/\text{l}$.

5. Sediments in Back Bay and within 500 meters of Giant Mine contain arsenic levels of 440 and 1300 mg/kg (milligrams per kilogram) respectively. Sediments taken from Back Bay at the mouth of Baker Creek contain arsenic concentrations of 2000 to 3000 mg/kg.
6. Arsenic levels in several Yellowknife area lakes exceed 0.1 mg/l.
7. Arsenic concentrations in snow usually range from 0.5 to 10 mg/l.
8. Soils in the City contain a highly variable amount of arsenic (1-600 ppm). High levels (> 4000 ppm) occur near Con and Giant Mine stacks.
9. Locally grown vegetables contain 0.02 to 7.0 ppm (wet weight) of arsenic.
10. Fish in Yellowknife Bay contain 0.2 to 0.3 ppm arsenic in their flesh.
11. Aquatic biota are almost non-existent in Baker Creek below Giant Mine. Benthic invertebrates are scarce in Back Bay.
12. Arsenic in small wildlife ranged from < 0.1 to 1.4 ppm.
13. Out of a total of 700 Yellowknife residents, 63 had arsenic levels > 10 ppm in hair. Of those 63 people, 57 received a more thorough examination, and 6 had arsenic levels exceeding 100 ppm.

14. Out of a total of 213 Cominco Con Mine employees, 160 had arsenic levels $\leq$ 20 ugm As/litre in their urine. Five employees had arsenic levels $\geq$ 100 ugm As/litre in their urine. Two of these five had lived in Yellowknife and worked at Con Mine for less than two years.

TABLE OF CONTENTS

	<u>Page No.</u>
Abstract	i
Table of Contents	iv
List of Figures	v
List of Tables	vi
1. Introduction	1
2. Summary of Technical Data	
2.1 Air	3
2.2 Water	13
2.3 Snow	29
2.4 Soil	35
2.5 Vegetation	46
2.6 Fish and Other Organisms	54
2.7 Wildlife	57
2.8 Human Hair and Urine	58
References	61

LIST OF FIGURES

		<u>Page No.</u>
Figure 1	Map of Yellowknife and Vicinity	2(A)
Figure 2	Map of Snow Sampling Stations, February 17, 1975, Yellowknife	30
Figure 3	Location of Soil Sample Sites, Con Mine Disturbed Soils Study, 1975	39
Figure 4	Location of Soil Sample Sites, Yellowknife Disturbed Soil Study, 1975	40
Figure 5	Location of Soil Sample Sites, Giant Mine Disturbed Soils Study, 1975	41
Figure 6	Road Dust Sample Sites, Yellowknife Disturbed Soils Study, 1975	43

LIST OF TABLES

	<u>Page No.</u>
1. Average Ore Processed and Arsenic Released to Atmosphere, Cominco Con Mine.	3
2. Average Ore Processed and Arsenic Released to Atmosphere, Giant Yellowknife Mine.	5
3. Summary of Arsenic Concentrations in Air, Various Communities.	7
4. Giant Yellowknife Mine Stack Arsenic Emission Tests	8
5. Ambient Air Survey, Cominco Con Mine, 1975-1976.	9
6. Continuous Air Sampling by Cominco Con Mine	11
7. Ambient Air Monitoring Program in the Yellowknife Area, 1975.	12
8. Criteria and Guidelines for Maximum Recommended Levels of Arsenic in Surface Water	13
9. Standard Limits for Arsenic in Water Supplies	14
10. Arsenic Concentration of Yellowknife Municipal Water Supply	15
11. Arsenic Concentration of Giant Yellowknife Mine - Mill Water Supply	18
12. Arsenic Content of Surface Water, Yellowknife, N.W.T., 1976.	19
13. Sediments, Yellowknife Bay, 1974.	20
14. Sediments, Yellowknife Bay, July-August, 1976. (Shoreline transects)	21
15. Sediments, Yellowknife Bay, July-August, 1976. (Transects out into Bay)	22
16. Sediments, Yellowknife Bay, July-August, 1976. (Near Niven Lagoon)	23
17. Arsenic Levels in Yellowknife Area Lakes	24
18. Occurrence of Arsenic in Various Water Bodies	27

	<u>Page No.</u>
19. Summary of Arsenic Deposition Rates, Yellowknife, N.W.T.	28
20. Arsenic Concentrations in Snow in Yellowknife Area	29
21. Spring Runoff Sampling Conditions, Yellowknife, N.W.T., 1976.	31
22. Spring Runoff Analyses, Yellowknife, N.W.T., 1976.	32
23. Yellowknife Arsenic Survey, 1970, (Soils) Cominco Ltd.	35
24. Yellowknife Arsenic Survey, 1975, (Soils) Cominco Ltd.	36
25. Yellowknife Arsenic Survey, 1976, (Soils) Cominco Ltd.	37
26. Arsenic Content of Soil Samples, Disturbed Soils Study, 1975.	38
27. Arsenic Content in Road Dust Samples, Disturbed Soils Study.	42
28. Various Soil Studies and Available Arsenic Data	44
29. Arsenic Concentrations in Yellowknife Area, 1970.	45
30. Allowable Arsenic Limits in Food, Food & Drug Act.	46
31. Arsenic Concentrations in Vegetables, Yellowknife, N.W.T.	47
32. Arsenic Content of Vegetation	49
33. Arsenic Content in Yellowknife Fruits and Vegetables Collected August, 1971.	50
34. Arsenic Content in Yellowknife Vegetables, September, 1976.	52

35. Arsenic Content of Native Grass Samples, Yellowknife, N.W.T.	53
36. Arsenic in Fish Livers and Fish Muscles, 1973.	54
37. Arsenic in Fish Muscle, 1975.	55
38. Organisms in Yellowknife Area Lakes and Rivers	56
39. Arsenic Content of Small Wildlife, Yellowknife Area, 1976.	57
40. Arsenic in Human Hair Samples, Yellowknife, 1975.	58
41. Arsenic in Human Urine Samples, Cominco Con Mine, Yellowknife, 1977.	59

1. INTRODUCTION

The Standing Committee on Arsenic Pollution in Yellowknife is comprised of members from the following government departments and agencies :

- (a) Environmental Protection Service (EPS),
Department of Fisheries and the
Environment (DFE),
- (b) Water Management Branch, Department of
Indian and Northern Affairs (DINA),
- (c) Mine Inspection Branch, DINA,
- (d) Medical Services, N.W.T. Region, National
Health & Welfare (NHW), Edmonton,
- (e) Northern Health Services, NHW, Yellowknife,
- (f) Health Care Plan, Government of the
Northwest Territories (GNWT),
- (g) City of Yellowknife,
- (h) Northwest Territories Chamber of Mines, and
- (i) Government of the Northwest Territories.

It is chaired by Mr. Dan Billing, Chief Environmental Protection Officer, GNWT.

At the request of this Standing Committee and its Chairman, a summary of data collected in the Yellowknife

Environmental Survey (1975) has been prepared.

This summary document is a compilation of tabulated data of arsenic concentrations in the various segments of the Yellowknife and vicinity environment. Air, water, snow, soil, vegetation, fish and wildlife samples were analysed during the Yellowknife Environmental Survey. Human hair sampling and analyses were undertaken by the Department of National Health and Welfare.

Included in this technical summary are data from the Food & Drug Act outlining allowable arsenic limits in food, standard limits for arsenic in water supplies, guidelines for allowable levels of arsenic in surface water, and the occurrence of arsenic in various water bodies. Standards on the arsenic level in soils in absence of a source of arsenic contamination, and available data on arsenic concentration in soils in other areas, have been tabulated. Arsenic concentrations in ambient air of some communities in North America are provided.

Yellowknife Bay Area

PERCENTAGE 0 100 200 300 400 500 600 700 800 900 1000
SCALE
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100




TABLE 1

Average Ore Processed and Arsenic
Released to AtmosphereCominco Con Mine,
Yellowknife, N.W.T.

(In pounds of arsenic per day)

Year	Ore Processed (tons/day)	Arsenic Removed from Waste Gases (lbs/day)	Arsenic Released to Atmosphere (lbs/day)
1948		-	2385
1949		2147	4161
1950		4043	459
1951		5117	494
1952		4969	310
1953		5926	313
1954		6630	409
1955		6954	430
1956		7114	420
1957		7870	401
1958	516	7353	384
1959	524	7302	431
1960	520	7502	599
1961	528	7129	712
1962	538	7429	595
1963	525	6898	484
1964	445	4070	222
1965	412	4940	370
1966	451	4366	311
1967	430	3600	316

TABLE 1 (cont'd)

Year	Ore Processed (tons/day)	Arsenic Removed from Waste Gases (lbs/day)	Arsenic Released to Atmosphere (lbs/day)
1968	413	3450	318
1969	400	2518	333
1970	396	2030	392
1971	425	0	0
1972	451	0	0
1973	462	0	0
1974	398	0	0
1975	404	0	0
1976		0	0
		21,780 tons	2,782 tons

(Data supplied by Cominco Con Mine)

TABLE 2

Average Ore Processed and Arsenic
Released to AtmosphereGiant Yellowknife Mine,
Yellowknife, N.W.T.

(In pounds of arsenic per day)

Year	Ore Processed (tons/day)	Arsenic to Underground Storage (lbs/day)	Arsenic to Tailings Pond (lbs/day)	Arsenic Removed by Hot Cottrell (lbs/day)	Arsenic to Stack (lbs/day)	Baghouse Efficiency (%)
1948	250					
1949	232					
1950	346					
1951	416				2053	
1952	850				3941	
1953	673				7094	
1954	756				11,980	
1955	788				6392	
1956	813				5998	
1957	848				6544	
1958	782				3330	
1959	989				115	
1960	991				165	
1961	1003				330	
1962	1029				330	

TABLE 2 (cont'd)

Year	Ore Processed (tons/day)	Arsenic to Underground Storage (lbs/day)	Arsenic to Tailings Pond (lbs/day)	Arsenic Removed by Hot Cottrell (lbs/day)	Arsenic to Stack (lbs/day)	Baghouse Efficiency (%)
1963	1063				330	
1964	1094		416		692	
1965	1082		509		372	
1966	1053	31,329	449	1014	537	98.3
1967	876	27,169	262	1168	287	99.0
1968	1024	30,941	54	576	502	98.8
1969	1095	30,552	56	551	661	97.7
1970	1164	28,793	76	588	485	98.3
1971	1106	26,393	91	701	1933	93.2
1972	1096	24,847	101	773	875	96.6
1973	1067	23,784	95	726	549	97.7
1974	899	19,643	99	728	485	97.6
1975	1074	19,338	106	858	479	97.6

6.

(Data supplied by Giant Yellowknife Mine)

TABLE 3

Summary of Arsenic Concentrations in Air

Various Communities

(Arsenic Concentration in $\mu\text{gm}/\text{m}^3$)

(Reference 1)

Location	1964		1972		1973		1974	
	Max.	Yearly Ave.	Max.	Yearly Ave.	Max.	Yearly Ave.	Max.	Yearly Ave.
Ontario:								
Windsor			0.300	0.030	0.025	0.012		
Welland			2.313	0.696	0.610	0.271	4.256	0.730
Sault Ste. Marie			0.040	0.017	0.057	0.013		
South Porcupine			--	--	1.090	0.168		
Sudbury			0.153	0.026	0.058	0.013		
Yellowknife, N.W.T.:								
Airport					0.62	0.12	1.46	0.12
Townsite					0.42	0.14	0.75	0.098
Giant Mine					0.95	0.22	1.68	0.26
United States:								
Chicago	0.04	0.03						
Toledo, Ohio	0.18	0.09						
El Paso, Texas	1.40	0.75						
Dallas, Texas	0.03	0.02						
Seattle, Washington	0.14	0.08						
Charleston, W.Virginia	0.36	0.25						
Average of 133 U.S. Stations								
		0.02						

TABLE 4

Giant Yellowknife Mine Stack
Arsenic Emission Tests
(Conducted by EPS, Edmonton staff)
(Reference 2)

Date	Test Number	Arsenic Emission (lbs As/day)
August 14, 1975	1	150.3
August 15, 1975	2	238.3
August 16, 1975	3	155.8
August 17, 1975	4	175.1
August 18, 1975	5	157.3
August 19, 1975	6	127.6
(average)		(167.4)

TABLE 5

Ambient Air Survey, Cominco Con Mine,
1975-1976.

Arsenic Concentration in the Air
Yellowknife, N.W.T.

(Arsenic Concentration in $\mu\text{g}/\text{m}^3$)

Reference 15

Sampling Period	Location	Total Arsenic Concentration (monthly geometric mean)
September 1975	North side of Con Mill	0.16
October 1975	" "	0.09
November 1975	" "	0.02
December 1975	" "	0.02
January 1976	North side of Con Mill	0.07
February 1976	" "	0.04
March 1976	" "	0.05
April 1976	" "	0.05
May 1976	" "	0.38
June 1976	June 18 - moved to Con Camp	0.12
July 1-28, 1976.	Con Camp	0.03
July 29 - August 24, 1976.		Test suspended
August 25 - Sept 1, 1976.	Con Camp	0.05
September 1976	Con Camp	0.05
October 1976	Con Camp	0.02
November 1976	Con Camp	0.04

Notes on Table 5

- (1) Annual geometric mean = 0.06 ug/m^3 arsenic
- (2) Test suspended July 29-August 24, 1976, due to labour dispute and staff shortage.
- (3) Samples obtained by Hi-Vol Sampler.
- (4) Chemical analysis at Cominco Laboratory, Trail, B.C.
- (5) British Columbia Government Control Branch objectives state that the annual geometric mean should not exceed 0.2 ug/m^3 arsenic.

TABLE 6

Continuous Air Sampling by Cominco Con Mine

Arsenic Concentration in the Air

Yellowknife, N.W.T.

(Arsenic Concentration in $\mu\text{g}/\text{m}^3$)

Sampling Date: 1976	Location	Arsenic Concentration
March 3-10	North side of Con Mill	0.05
March 10-17	" " " "	0.07
March 17-24	" " " "	0.03
March 24-31	" " " "	0.07
March 31- April 7	" " " "	0.03
April 7-14	" " " "	0.06
April 14-21	" " " "	0.08
April 21-28	" " " "	0.04

Note: (1) Samples obtained by Hi-Vol Sampler

(2) Chemical analysis at Cominco Laboratory,
Trail, B.C.

TABLE 7

Ambient Air Monitoring Program

in the Yellowknife Area, 1975.

Arsenic Concentration in Suspended Particulates
(Reference 3)

Parameter	Date	Arsenic Concentration
Overall annual geometric mean suspended particulate	1973	0.08 $\mu\text{gms As./m}^3$
	1974	0.09 "
	1975	0.06 "
Individual 24-hour suspended particulate	1973-1975	<0.01 - 3.91 $\mu\text{gms As./m}^3$ range

Following are available data on standards for arsenic in air.

Extraneous Air Quality Limits: ($\mu\text{gm As./m}^3$)

Arsenic Concentrations in
Suspended Particulates: 24-hour basis

B.C. Objective	-	1.0
USSR Standard	-	3.0
Czechoslovakia Standard	-	3.0

TABLE 8

Criteria and Guidelines for
Maximum Recommended Levels of Arsenic
in Surface Water

(Arsenic Concentration in ppm)

	Water Quality Criteria U.S. Environmental Protection Agency	Maximum Arsenic Level: NHW (1968) Guidelines
Public Water Supplies (raw water from rivers or lakes used for drinking; i.e. casual water sources)	[*] ≤ 0.1 recommended	0.05 maximum acceptable 0.01 maximum desirable
Recreation and Esthetics	not specified	not specified
Fresh Water Aquatic Life & Wildlife	not specified	not specified
Marine Aquatic Life and Wildlife	0.01 and 0.05	not specified
Agricultural Uses of Water	0.01 (continuous use)	
Water for livestock	2.0 (for up to 20 years)	0.05
Water for irrigation	not specified	2.0 maximum acceptable 0.1 maximum desirable
Industrial Water Supplies	not specified	not specified

* 0.1 mg/l (ppm) total arsenic. Obtained from Water Quality Criteria 1972, p. 56; EPA. R3.73.033 March 1973, A Report of the Committee on Water Quality Criteria, Environmental Studies Board, National Academy of Sciences, Washington, D.C.

TABLE 9

Standard Limits for Arsenic in Water Supplies

(Arsenic concentration in ppm.)

1.	U.S.A. Public Health Service (1962)	
	a) Limit which should not be exceeded if an alternative water source can be made available	0.01
	b) Limit above which water source should be rejected	0.05
2.	Canada, Department of National Health & Welfare (1968)	
	a) Acceptable Limit	0.01
	b) Maximum permissible limit	0.05
3.	World Health Organization	
	International standard (1971)	0.05
4.	World Health Organization	
	European standard (1970)	0.05

TABLE 10

Arsenic Concentration of
Yellowknife Municipal Water Supply

(Arsenic Concentration in ppm.)

Date of Sampling	Tap Water	Schooldraw Pumphouse Yk. River Supply	Schooldraw Pumphouse Yk. Bay Supply
<u>1975 Summary</u>	0.01	0.01	
Feb. 18, 1975			0.011
June 18, 1975			0.020
Sept. 28, 1975			0.01
Oct. 28, 1975			0.01
Nov. - Dec., 1975			0.01
<u>1976:</u>			
January 6	<0.01		
January 13	<0.01		
January 19	<0.01		
January 26	<0.01		
February 9	0.0008		
February 11		0.0008	0.003
March 2	0.0029		
March 9	0.0006		
March 16	0.0241		
March 22	0.003		
March 30	0.0006		
April 5	0.0008		
April 13		0.0007	0.0045
April 27		0.0001	0.0009
May 11		0.0005	0.020
May 17	0.0020		
May 25	0.0068		
June 7	<0.0005		
June 11		0.0046	0.014
June 14	0.0010		
June 21	0.0008		

TABLE 10 (cont'd)

Date of Sampling	Tap Water	Schooldraw Pumphouse Yk. River Supply	Schooldraw Pumphouse Yk. Bay Supply
<u>1976:</u>			
July 5	0.0006		
July 12	0.0038		
July 19	0.0020		
July 29	0.0001		
August 2	0.0002		
August 10	<0.0005		
August 13		0.002	0.55
August 23	<0.0005		
August 30	0.0008		
September 14	<0.0005		
September 23	<0.0005		
September 27	<0.0005		
October 5	0.0018		
October 15		<0.0005	0.014
October 18	<0.0005		
November 8	<0.0005		
November 15	<0.0005		
November 29	<0.0005		
December 7	<0.0005		
December 14	<0.0005		
<u>1977:</u>			
January 4	<0.0005		
January 6		<0.0005	0.0093
January 11	0.0007		
January 18	0.0008		
January 24	0.0010		
January 31	<0.0005		
February 8	<0.0005		
February 14	<0.0005		
February 22	<0.0005		
March 2	<0.0005		

TABLE 10 (cont'd)

Date of Sampling	Tap Water	Schooldraw Pumphouse Yk. River Supply	Schooldraw Pumphouse Yk. Bay Supply
<u>1977:</u>			
March 7	< 0.0005		
March 14	< 0.0005		
March 23	< 0.0005		
March 28	< 0.0005		
April 4	< 0.0005		
April 20	0.0006		

TABLE 11

Arsenic Concentration of
Giant Yellowknife Mine
Mill Potable Water Supply

Year	Month	Ave. Arsenic Content ppm.	Minimum ppm.	Maximum ppm.
1973	January	0.0077	N.D.	0.014
	February	0.0066	0.001	0.013
	March	0.0069	0.002	0.013
	April	0.0098	0.005	0.015
	May	0.0260	0.006	0.144
	June	0.0460	0.005	0.290
	July	0.0300	0.004	0.097
	August	0.0210	0.007	0.045
	September	0.0195	0.011	0.037
	October	--	--	--
	November	0.0220	N.D.	0.056
	December	0.0200	0.000	0.103
1974	January	0.0110	0.000	0.040
	February	--	--	--
	March	0.0120	0.003	0.024
	April	0.0100	0.005	0.019
	May	0.0090	0.005	0.016
	June	0.0070	0.005	0.013
	July	0.0270	0.003	0.093
	August	0.0190	0.005	0.048
	September	--	--	--
	October	0.0097	0.005	0.019
	November	0.0080	0.003	0.013
	December	0.0070	0.003	0.011
1975	January	0.0120	0.005	0.019
	February	0.0080	0.003	0.011
	March	0.0080	0.005	0.013
	April	0.0160	N.D.	0.035
	May	0.0470	<0.015	0.190
	June	0.0480	0.021	0.320
	July	0.0410	<0.015	0.186
	August	--	--	--
	September	0.0320	<0.015	0.073
	October	0.0260	<0.015	0.057
	November	--	--	--
	December	0.0200	<0.015	0.060
1976	January	0.0170	<0.015	0.059
	February	0.016	<0.015	0.027
	March	0.017	<0.015	0.042
	April	0.024	<0.015	0.058
	May	0.033	<0.015	0.100
	June	0.056	<0.015	0.189

N.D. = Not Determined

TABLE 12

Arsenic Content of Surface Water

Yellowknife, N.W.T. (1976)

(EPS, Yellowknife)

BAKER CREEK - above mine

	<u>May 26</u>	<u>June 10</u>	<u>June 24</u>	<u>July 7</u>	<u>July 21</u>	<u>Aug. 12</u>	<u>Aug. 24</u>	<u>Sept. 3</u>
mg/l	0.03	<0.02	<0.02	<0.02	0.10	<0.02	<0.02	0.03

YELLOWKNIFE BAY

Station 1

	<u>May 17</u>	<u>May 28</u>	<u>June 1</u>	<u>June 3</u>	<u>June 9</u>	<u>June 23</u>	<u>July 8</u>	<u>July 26</u>	<u>Aug. 5</u>	<u>Aug. 18</u>	<u>Oct. 6</u>
<0.02	<0.02	<0.02	0.74 0.03 <0.02	0.05 <0.02 <0.02	0.06	0.40	<0.02	<0.02	<0.02	<0.02	<0.02

Station 2

0.03	<0.02	No data	No data	<0.02	0.04	0.04	<0.02	<0.02	<0.02	<0.02
------	-------	---------	---------	-------	------	------	-------	-------	-------	-------

Station 3

<0.02	0.02	"	"	0.03	0.03	0.11	<0.02	<0.02	<0.02	<0.02
-------	------	---	---	------	------	------	-------	-------	-------	-------

Station 4

0.03	No data	"	"	No data	No data	No data	<0.02	<0.02	<0.02	<0.02
------	---------	---	---	---------	---------	---------	-------	-------	-------	-------

YELLOWKNIFE RIVER

	<u>May 5</u>	<u>May 26</u>	<u>June 10</u>	<u>June 24</u>	<u>July 7</u>	<u>July 21</u>	<u>Aug. 24</u>	<u>Sept. 1</u>
<0.02	<0.02	<0.02	<0.02	<0.02	0.03	0.08	<0.02	<0.02

TABLE 13

Sediments, Yellowknife Bay (1974)

Yellowknife, N.W.T.

(EPS, Yellowknife)

(readings in mg/kg, dry weight)

a)	<u>Near Yellowknife River</u>			
	<u>Mean</u>	<u>Range</u>	<u>95% Confidence Limit</u>	<u>Number of Samples</u>
	138	6-515	83	14
b)	<u>Back Bay</u>			
	441	33-1332	254	11
c)	<u>0-500m from Mine</u>			
	1316	385-2400	480	6
d)	<u>Near Latham and Jolliffe Island</u>			
	130	34-427	75	11
e)	<u>Between Con Mine and Detah</u>			
	63	8-227	47	9
f)	<u>Wool Bay</u>			
	11	10-12	--	3

TABLE 14

Sediments, Yellowknife Bay (July-August, 1976)
Yellowknife, N.W.T.

(EPS, Yellowknife)

(Readings in mg Arsenic/kg, dry weight (ppm))

Distance from Point of Waste Disposal at Mouth of Baker Creek (meters)	Arsenic Level in Sediments	
	Transect # 1 (N)	Transect # 2 (S)
0	2050	2050
200	325	775
400	120	550
600	120	825
800	140	300
1000	-	50

- Notes:
- (1) Distance increases from point of waste disposal at mouth of Baker Creek along the shoreline.
 - (2) Transect # 1 proceeds in a north direction from point of disposal.
 - (3) Transect # 2 proceeds in a south direction from point of disposal.
 - (4) Analyses carried out by Barringer Research (Toronto) using plasma emission.

TABLE 15

Sediments, Yellowknife Bay (July-August 1976)
Yellowknife, N.W.T.

(EPS, Yellowknife)

(Readings in mg Arsenic/kg, dry weight (ppm))

Distance from point of Waste Disposal at Mouth of Baker Creek (meters)	Arsenic Levels in Sediments		
	Transect # 1 (NE)	Transect # 2 (E)	Transect # 3 (SE)
0	3000	3000	3000
400	750	575	1450
800	475	185	1100
1200	75	425	20
1600	28	2400	-
2000	12	-	-
2400	12	-	-

Notes: (1) Transects proceed out NE, E & SE from disposal points.

(2) Analyses carried out by Barringer Research (Toronto) using plasma emission.

TABLE 16

Sediments, Yellowknife Bay (July-August 1976)
Yellowknife, N.W.T.

EPS, Yellowknife.

(Readings in mg Arsenic/kg, dry weight (ppm))

Distance from Niven Lagoon sewage outfall point at Back Bay (meters)	Arsenic Level
100	80
300	15
500	120

- Notes: (1) Distance is measured from sewage discharge point out into Back Bay.
- (2) Analyses carried out by Barringer Research (Toronto) using plasma emission.

TABLE 17

Arsenic Levels in Yellowknife Area Lakes

(arsenic concentration in ppm)

Lake/Location	Arsenic (soluble)	Remarks
"Lake 2; 1.9 miles S.W."	3.90	Data from O'Toole
"Lake 7; 0.3 miles S.W."	12.40	Reference 4
"Lake 9; 1.9 miles S.W."	0.19	Summer, 1970
"Lake 13; 1.8 miles N.W."	0.56	
"Lake ; 10 miles"	0.20	
"Lake ; 20 miles"	0.001	
Yellowknife Bay:		
near Latham Island	0.010	Data from Cominco Ltd.,
near Mosher Island	<0.005	survey (1970). Wadey, R.H.
near Detah village	<0.005	Supervisor, Waste Control,
3000 ft. from Kam Point	<0.005	Cominco, Trail, B.C.
Stock Lake	0.005	
Frame Lake	0.055	
Pud Lake (discharge)	0.030	
Kam Lake	3.0	
Yellowknife Bay:		
near Latham Island	0.020	Cominco, Ltd. survey
near Mosher Island	0.022	(1971)
near Detah Village	0.005	
3000 ft. from Kam Point	0.029	
Stock Lake	0.130	
Frame Lake	0.250	
Pud Lake (discharge)	5.600	
Kam Lake	2.800	
Kam Lake	1.0 - 5.0	Data from memo Brunskill
Grace Lake	0.001- 0.10	to Hamilton Feb. 6/75
Frame Lake	0.150- 0.180	"Arsenic in Yk. Waters"
Lakes 16 miles N.E.	n/d	(Tests carried out over
Lakes 12-15 miles N. & N.E.	0.0005 - 0.007	1972-1974()n/d=non detectable)

TABLE 17 (cont'd)

Lake/Location	Arsenic (soluble)	Remarks
Long Lake	3.0 - 8.0	G. Brunskill; preliminary findings, 1975
Long Lake - beach	0.135	Feb. 25/75; G. Brunskill
Long Lake - center	0.105	"
Stock Lake	0.112	"
Range Lake	1.296	Feb. 26/75; G. Brunskill
Fault Lake	0.270	"
Rat Lake	0.471	"
Kam Lake	2.850	"
Frame Lake	0.562	Feb. 7/75; G. Brunskill
Grace Lake	0.026	"
Yellowknife Bay:		
near Latham Island	0.01	Cominco, Ltd. survey (1975)
near Mosher Island	0.01	Brown, R.L., Supervisor,
near Detah Village	0.01	Waste Control, Cominco,
3000 ft. from Kam Point	0.01	Trail, B.C.
Arden Bay	0.01	
Stock Lake	0.06	
Frame Lake	0.21	
Kam Lake	2.55	
Long Lake	0.09	
Kam Lake:		
North basin	2.04	Data collected May 25/76
Central	1.08	by Dr. J. Moore, EPS,
South basin	2.06	Yellowknife. Internal correspondence.

Notes: (1) Dr. G. Brunskill's analyses (acid digestion method) carried out at the Freshwater Institute, Winnipeg, Manitoba.

(2) Dr. J. Moore's analyses carried out at the E.P.S. laboratory, Edmonton, Alberta.

TABLE 17 (cont'd)

Lake/Location	Arsenic (soluble)	Remarks
Yellowknife Bay:		
near Latham Island	0.04	Cominco Ltd Survey (1976)
near Mosher Island	0.04	(Reference 15)
near Detah Village	0.01	
near Kam Point	0.005	
Arden Bay	0.01	
Long Lake	0.14	
Stock Lake	0.15	
Frame Lake	0.33	
Kam Lake	1.70	

Notes: (cont'd)

- (3) Cominco analyses carried out at Trail, B.C.
(arsine generation method)

TABLE 18

Occurrence of Arsenic in Various Water Bodies

Arsenic in Water (conc. in ppm)

(Reference 1)

Occurrence	Level or Range of Levels
Sea Water	0.006 - 0.30
River water, United States	mean 0.0004; maximum 0.230
Natural levels, Canadian rivers; 1968 - 1974	0.005 - 0.13
Lakes Superior, Ontario, Huron, Erie	0.00025 - 0.001
River; Sweden (upstream from industrial site)	0.002
Surface water; United States	mean 0.064; range 0.005 - 0.336
Tap Water; United States	0.100 in some samples tested
Spring water - California, Rumania USSR; New Zealand	0.400 - 1.300 (high in bicarbonate)
Near Yellowknife, 1973 - 1974	0.070 - 1.00
Great Slave Lake	0.004 - 0.500 total 0.001 - 0.030 dissolved
Taiwan (artesian wells)	0.010 - 1.82 (most 0.40 - 0.60)
Cobalt Lake (Ottawa River Basin) 1970 - 1971	1.00 - 2.50
Argentina, Cordoba province	approximately 1.50
New Zealand (fresh water)	approximately 20.0
Searles Lake, California (high salinity water)	over 200.0

TABLE 19

Summary of Arsenic Deposition
Rates, Yellowknife, N.W.T.

Survey Date	Survey	Arsenic Deposition (lbs/mile ² /month)		Details of Survey
1. 1950-1963	de Villiers & Baker (Ref. 5)	low	- 17	Survey in town of Yellowknife
		high	- 469	
		average	- 106	
2. 1950-1956	Cominco Con Mine	low	- 27	6 sample points: ½ to 2 miles from stack
		high	- 267	
3. 1955-1956	Giant Mine	low	- 27	15 sample points: 1, 2 and 3 miles from stack
		high	- 613	
4. 1974-1975	E.P.S. Snow Survey (Reference 6)	low	- 1.2	52 sample points: 'low' was obtained 4 miles south of Giant; 'high' was obtained inside Giant property. (Analysis by Vasak/ Sedevic colorimetric method.)
		high	- 564	
		calculated winter mean	- 11	
5. 1975	E.P.S. Ambient Air (Ref. 3)	low	- 2.2	'average' is based on results of 22 stations over period June to October 1975. (Analysis by Vasak/Sedevic colorimetric method and by XRF.)
		high	- 37	
		average	- 9.8	

- Notes: (a) Surveys 1, 2, 3 & 5 are based upon results of fall-pan surveys.
- (b) Survey 4 is based upon results of snow coring.
- (c) Samples from EPS surveys were analysed at EPS Laboratory, Edmonton, Alberta.

TABLE 20

Arsenic Concentrations in Snow in Yellowknife Area

(Arsenic Conc. in ppm.)

Survey	Sample	Location	Arsenic	Acidity (pH)	Remarks
O'Toole (Reference 4)	'Snow 5'	0.35 mile N.E.	0.68		
	'Snow 9'	1.30 mile N.E.	1.20		
	'Snow 10'	0.22 mile N.W.	8.75		
	'Snow 21'	2.0 miles N.E.	0.47		
EPS, Yellowknife district office (Feb. 17, 1975)	1	(Sample locations on attached map; figure 2).	0.46	3.4	Analyses carried out at the Water Quality Laboratory, D.F.E., Calgary, Alberta.
	2		0.44	3.7	
	3		0.48	3.9	
	4		0.81	3.6	
	5		11.40	6.6	
	6		0.71	6.7	
	7		9.10	4.4	
	8		1.30	4.4	
	9		8.60	4.2	
	10		0.50	4.2	
	11		0.027	4.2	
	12		2.30	3.6	
March, 1975 survey (6)	(Refer to report, reference # 6)	0.1 mile W. of Giant	0.17	4.6	average; 52 samples. maximum minimum
			8.80	7.7	
			0.02	2.7	

Yellowknife Bay Area




FIGURE 2 MAP OF SNOW SAMPLING STATIONS; FEBRUARY 17, 1975, ENVIRONMENTAL PROTECTION SERVICE YELLOWKNIFE, NWT.

TABLE 21

Spring Runoff Sampling Conditions
Yellowknife, N.W.T.

Sampling Date	Conditions
1) April 13, 1976	Spring melt in the early stage; minimal flows; ice cover on all lakes and rivers.
2) April 27, 1976	Melting rate near its maximum; Yellowknife River was flowing; surface runoff flows peaking; most lakes were melting around edges.
3) May 11, 1976	Melt was nearing completion; surface runoff was beginning to decrease; most lakes had only patches of ice remaining.
4) June 11, 1976	Several drainage ditches and surface melt; water pools were dry.

TABLE 22

1976 Spring Runoff Analyses

Yellowknife, N.W.T.

(Arsenic Concentration, ppm.)

Sampling Site	Arsenic Concentration, ppm.				Remarks
	1 April 13	2 April 27	3 May 11	4 June 11	
Yellowknife River	0.09	0.001	0.0009	0.008	At bridge on Ingraham Trail; (April 13, meltwater on ice surface)
Giant Tailings Pond	13.9	2.8	5.7	6.0	April 13 - meltwater at edge of tailings area; 2, 3 & 4 - seepage from pond.
Giant Surface Meltwater	0.023	0.23	0.30	dry	Shallow pool near road at tailings pond.
Giant Tailings Effluent	31.0	4.20	14.4	9.5	Entering Baker Creek.
Baker Creek	1.20	1.20	0.70	0.80	Represent As levels after tailings effluent diluted with Baker Creek.
Mouth of Baker Creek	1.50	1.00	0.73	1.32	Discharge to Yellowknife Bay.
Junction of Giant Road and Airport Road	0.036	0.28	0.068	0.17	Slough on N.E. side.
Runoff to Stock Lake	0.075	0.184	0.160	0.096	1 - pool near Bristol monument; 2 & 3 - flow into Stock Lake; 4 - Stock Lake water.

TABLE 22 (cont'd)

(Arsenic Concentration, ppm.)

Sampling Site	1 April 13	2 April 27	3 May 11	4 June 11	Remarks
Long Lake	0.030	0.048	0.074	0.122	1 - near boat ramp/picnic area; 2 & 3 - edge of lake; 4 - in lake.
Surface Meltwater near Northland Trailer Court	0.014	0.056	0.035	1.45	1, 2, 3 - S.E. side of junction Franklin Avenue & Correctional Institute road; 4 - Kam Lake.
Yellowknife Bay at Con Mine Pumphouse	0.032	0.015	0.016	0.22	1 & 4 - inside pumphouse; 2 & 3 - from bay outside pump- house; (water used only for industrial purposes).
Back Bay/Rainbow Valley	0.177	0.08	0.016	0.36	1 - meltwater on surface of ice; 2 & 3 - near shoreline.
Rat Lake	0.16	0.45	0.40	0.56	1, 2 & 3 - small stream entering Rat Lake; 4 - from Rat Lake (stream dry).
Con Mine Tailings	4.0	4.2	7.4	3.6	1 - meltwater near dam in Pud Lake; 2 & 3 - control dam outflow; 4 - inside tailings pond.
Meltwater near City Snow dump area	0.044	0.056	0.035	0.058	From ditch near snow dump; ½ mile N. of Niven Lake; east side of highway.

TABLE 22 (cont'd)

Sampling Site	(Arsenic Concentration, ppm.)				Remarks
	1 April 13	2 April 27	3 May 11	4 June 11	
Yellowknife Bay at Schooldraw Culvert	0.012	0.052	0.024	0.067	Runoff under Schooldraw Road into Yellowknife Bay.
Detah Water Hole	0.0043	0.0070	0.0050	0.0034	Just offshore from Detah Village.
Yellowknife Water Supply	0.0007	0.0001	0.0005	0.0046	Schooldraw Road pumphouse; water from Yellowknife River.
Yellowknife Emergency Water Supply	0.0045	0.0090	0.020	0.014	Yellowknife Bay; wetwell in schooldraw pumphouse.
Causeway to Latham Island	0.0036	0.007	0.017	0.036	Northside of causeway.
Yellowknife tapwater	0.0005	0.0008	0.0005	0.0005	Tapwater normally sampled weekly. Canadian Drinking Water Standard - 0.01 ppm.
McNiven Beach slough	---	0.012	0.015	0.011 (0.27)	1 - frozen solid, no sample; 2 & 3 - slough draining to Frame Lake; 4 - slough out- flow minimal (Frame Lake sampled also).

TABLE 23

Yellowknife Arsenic Survey, 1970 (Soils)

Cominco, Ltd., Trail, B.C.

(Arsenic Concentration in dry weight, ppm.)

Location	Depth in Soil, Inches				
	0-1	1-7	7-13	0-6	0-13
Undisturbed Soil:					
1) 0.75 mile S.W. of Giant stack.				58	
2) 2 miles S. of Giant stack.	205	50	14		
3) East end, Frame Lake.	101	45	26		
4) Con residential area.	147	33			
5) East of Negus mine.				96	
6) West of Con head-frame.				2980	
Cultivated Soil:					
7) Garden, Con residential area.				21	40

Standard

Arsenic level in soils in the
absence of a source of contamination... ≤ 5.0 ppm.

- Notes: (1) Analyses in Tables 23, 24 and 25 conducted at Trail Laboratory.
- (2) Surveys carried out in 1975 and 1976 are reported in Reference 15.

TABLE 24

Yellowknife Arsenic Survey, 1975 (Soils)

Cominco Ltd, Trail, B.C.

(Arsenic Concentration in dry weight, ppm)

Location	Depth in Soil, Inches.			
	0 - 1	1 - 7	7 - 13	0 - 13
Undisturbed Soils:				
1. Con Property (100 yards N. of Robertson head- frame)	2880	380	21	
2. Negus Mine (100 yards N.E. of old shaft)	3680	4280	3600	
3. Rat Lake (25 yards W. of lake)	368	46	68	
4. Frame Lake (50 yards N. of lake)	68	92	12	
5. Giant - Edmonton road junction (75 yards N. of junction)	74	33	24	
6. Giant Mine (1 mile S. of stack)	392	157	40	
Cultivated Soil:				
7. Con Garden (D. White)				64
8. Town Garden (D. Egli) - N. end				24
S. end				21

TABLE 25

Yellowknife Arsenic Survey, 1976 (Soils)

Cominco Ltd, Trail, B.C.

(Arsenic Concentration in dry weight, ppm)

Location	Depth in Soil, Inches.			
	0 - 1	1 - 7	7 - 13	0 - 13
Undisturbed Soil:				
1. Con Property (100 yards N. of Robertson head- frame)	420	30	30	
2. Negus Mine (100 yards N.E. of old shaft)	1210	250	60	
3. Rat Lake (25 yards W. of lake)	20	20	40	
4. Frame Lake (50 yards N. of lake)	140	20	80	
5. Giant - Edmonton road junction (75 yards N. of junction)	30	40	40	
6. Giant Mine (1 mile S. of stack)	140	110	20	
Cultivated Soil:				
7. Con Garden (D. White)				80
8. Town Garden (D. Egli) - N. end				30
S. end				40

TABLE 26

Arsenic Content of Soil Samples, Disturbed Soils Study, 1975
(Reference 7)

Sample	Location	Total Arsenic (ppm)	Insoluble Arsenic (ppm)
M-1	Con Mine	192	191
M-2	Con Mine	342	341
M-3	Con Mine	58	57
M-4	Con Mine	115	114
M-5	Con Mine	115	115
M-6	Con Mine	490	484
S-1	Town	36	25
S-2	Town	27	8
S-3	Town	39	27
S-4	Town	10	10
S-5	Town	75	47
S-6	Town	138	138
S-7	Town	19	4
S-8	Town	10	5
S-9	Town	605	46
S-10	Town	23	17
S-11	Town	83	81
S-12	Town	43	23
S-13	Town	1	1
S-14	Town	26	1
S-15	Town	42	42
S-16	Town	8	1
S-17	Town	18	8
S-18	Town	21	7
S-19	Town	5	4
S-20	Town	5	5
S-21	Con Mine	313	313
S-22	Con Mine	129	129
S-23	Con Mine	755	755
S-24	Con Mine	10	10
S-25	Con Mine	605	605
S-26	Giant Mine	261	105
S-27	Giant Mine	38	36
S-28	Giant Mine	33	33
S-29	Giant Mine	79	70
S-30	Giant Mine	75	66




FIGURE 3 LOCATION OF SOIL SAMPLE SITES; CON MINE
DISTURBED SOILS STUDY; 1975




FIGURE 4 SOIL SAMPLE SITES; YELLOWKNIFE, NWT.
DISTURBED SOILS STUDY; 1975




FIGURE 5 SOIL SAMPLE SITES ; GIANT MINE
DISTURBED SOILS STUDY; 1975

TABLE 27

Arsenic Content in Road Dust Samples, Disturbed Soils Study.
(Reference 7)

Gravel Samples			Dust from Pavement		
Sample	Total Arsenic (ppm)	Insoluble Arsenic (ppm)	Sample	Total Arsenic (ppm)	Insoluble Arsenic (ppm)
G-1	74	74	P-1	33	21
G-2	31	25	P-2	128	128
G-3	30	30	P-3	58	58
G-4	21	20	P-4	216	206
G-5	38	38	P-5	94	94
G-6	49	49	P-6	58	58
G-7	29	28	P-7	161	155
G-8	21	21	P-8	38	38
G-9	16	16	P-9	45	45
G-10	42	34	P-10	36	31

42.

Notes on Tables 26 & 27

- (1) Soil samples M-1 to M-6 (Table 26) were analysed by 'acid digestion' at Chemical & Geological Laboratories, Edmonton.
- (2) Soil samples S-1 to S-30 (Table 26) were analysed by XRF at Northern Environmental Consultants, Edmonton.
- (3) Gravel and dust samples (Table 27) were analysed by XRF at Northern Environmental Consultants, Edmonton.




FIGURE 6 ROAD DUST SAMPLE SITES ; YELLOWKNIFE
DISTURBED SOIL STUDY; 1975

TABLE 28

Various Soil Studies and Available Arsenic Data

Study	Date	Location	Arsenic (ppm)	Remarks
O'Toole (4)	Summer, 1970	Yellowknife area	low: 34 high: 7598	Concentrations for the various samples are presented in the following table.
Temple (8)	1974	Ontario	107 35 10	Average of over 50 samples near smelter 'A'. Average of over 50 samples near smelter 'B'. Urban area, remote from smelters.
Cominco (9)	July, 1975	Yellowknife area	low: 12 high: 4280	Nine sample locations; 4 soil depths; outlined in Cominco report.
Toft (1)		Agricultural soils	0.1 to 40 range	Health Protection Branch report.
Toft (1)		Orchard soils	100	Health Protection Branch report.
Plambeck & Smith (10)	June - Sept. 1975	Yellowknife - Con Yellowknife - town Yellowknife - town Yellowknife - Giant	82 52 36 287	Garden 'A' Garden 'B' Garden 'C' Garden 'D'

TABLE 29

Arsenic Concentrations in Yellowknife Area

O'Toole, Clark, Malaby and Trauger

1970
(Reference 4)

Sample	Distance from Smelter (miles)	As (ppm)	Remarks
1	2.7 S.W.	219	Samples 1, 2, 4, 5, 6 and 7 are mineralized type soils
1A		199	
2	1.9 S.W.	115	Samples 1A, 2A, 4A, 5A, 6A, 7B are of high organic, mossy-type soils.
2A		164	
4	1.2 S.E.	601	
4A		814	
5	0.4 N.E.	602	
5A		852	
6	0.3 N.E.	3108	
6A		2641	
7	0.2 S.W.	6852	
7A		2449	
7B		7598	
8	0.8 N.E.	269	
9	1.3 S.W.	153	
12	1.8 N.W.	163	
13	1.8 N.W.	110	
14	0.7 N.W.	45	
TOWNSITES		119	
		34	
		178	

Garden soil samples from Yellowknife

TABLE 30

Allowable Arsenic Limits in Food

Food and Drug Act Regulations
 Health Protection Branch
 Department of National Health
 and Welfare

Food	Arsenic ppm
Marine & Freshwater Animal Products	5.0
Liver	1.0
Fresh Fruits	2.0
Fresh Vegetables	1.0
Apple Juice, Cider, Wine, Beer.	0.2
Other Fruit Juices, except Apple Juice	0.1
Beverages, as consumed, and Bottled Water	0.1
Tea	1.0
Fish Protein	3.5
Baking Powder	2.0
Gelatin	2.0
Dried Herbs and Spices	5.0

TABLE 31

ARSENIC CONCENTRATIONS IN VEGETABLES (ppm, natural weight)

Summary of Surveys in Yellowknife Area, including
Location of Garden, Test Laboratory and Method.

Vegetable	Con Mine Residences		Yellowknife Residences		Giant Mine Residences	
	Food & Drug Lab Ottawa 1970	Cominco Lab, Trail, B.C. 1970 1975	Food & Drug Lab Ottawa 1971 1975	Cominco Lab, Trail, Edmonton 1970 1975 (XRF)	Food & Drug Lab Ottawa 1971 1975	Cominco Lab, Trail, Edmonton 1975 1975 (XRF)
lettuce	2.65 2.00 1.32 1.07 1.42 1.55	0.64 1.28 0.26 0.28	2.25 1.10	0.28 46	0.16	59
chubarb		0.05 0.10 0.13	0.12 0.04	0.13	0.25	16
carrots		0.05 0.45 0.17	0.39 0.07 0.11 0.07	0.17 ≤ 10	0.22	
potatoes		0.20 0.03 0.31 0.25	0.07 0.02 0.08 0.06	0.25 16 ≤ 10	0.24	≤ 10
plant top)		(24) (19)				
beets		0.42 0.09 0.20 0.26	0.51 0.27	0.61 0.26	0.36	

TABLE 31 (cont'd)

Vegetable	Con Mine Residences		Yellowknife Residences		Giant Mine Residences	
	Food & Drug Lab Ottawa 1970	Cominco Lab, Trail, B.C. 1970 1975	Plambeck & Smith Edmonton 1975 (XRF)	Food & Drug Lab Ottawa 1971 1975	Cominco Lab, Trail, 1970 1975	Plambeck & Smith Edmonton 1975 (XRF)
Turnips		0.16	0.27			
Swiss Chard	1.05					
	3.15					
Small Carrots (leaves & roots)	4.15			2.20		
	3.10			1.15		
	1.15					
	1.36					
Beets (Leaves)	0.52					
Kohlrabi				0.67		
Cabbage				0.48		
Green Pea Pods				0.14		
Peas & Pods				0.08		
				0.11		
				0.06		
Tomatoes				0.13		
Onions				0.49		
				0.77		
Green Beans				0.25		
Plant tops			24			
Gooseberries			21			

TABLE 32

Arsenic Content of Vegetation (ppm.)

Vegetable	Fruit	Grass, Shrubs	Remarks (reference)
1.0	2.0		Federal maximum acceptable limits.
0.1		12-138 range	Level for vegetables grown in uncontaminated areas. O'Toole (4) 1970
<0.02	0.02-0.10		National Health and Welfare data (1).
	0.0 -5.0		Canadian apples, 1962-1964; higher results possibly due to use of arsenic compounds as pesticides on fruit crops.
<1.0	<1.0	1.0-62.0	Temple (8). Near smelters (1974).
<u><10-59 range</u>	<u><10-21 range</u>		Flambeck and Smith (10); 10 ppm was the detection limit of which the investigators were confident given the apparatus and technique used (XRF).

TABLE 33

Arsenic Content in Yellowknife Fruits and Vegetables Collected August, 1971
(Reference 1)

	Location	Arsenic (ppm) As Received	Arsenic (ppm) Washed & Dried	Arsenic (ppm) Cleaned as in Kitchen
1.	High Bush Cranberry	Con Mine	0.15	
2.	Gooseberry	Con Mine	0.48	0.19
3.	Black Currant	Con Mine	0.63	0.18
4.	Low Bush Cranberry	Airport	0.32	
5.	High Bush Cranberry	Airport	0.43	
6.	Vegetables, leaves	Mr. Bugg (204B; Giant Mine)	6.80	2.7
7.	Vegetables, leaves	Mrs. Richardson-Con Mine	1.70	0.53
8.	Vegetables, leaves	Mr. A.P. Morris	1.40	0.54
9.	Vegetables, leaves	Correctional Camp	0.32	0.12
10.	Vegetables, leaves	Mr. Christensen-Old Town	1.10	0.41
11.	Vegetables, leaves	5018-54 Street, Yellowknife	0.30	0.10
12.	Carrots	Mrs. Bugg (204B; Giant Mine)	0.61	0.39
	Beets	"	1.60	0.51
	Rhubarb	"		0.12
	Potatoes	"	0.96	0.07
13.	Carrots	Mrs. Richardson-Con Mine	0.25	0.07
	Beets	"	0.49	0.27
	Peas and Pods	"	0.16	0.08
	Potatoes	"	0.45	0.02

TABLE 33 (cont'd)

	Location	Arsenic (ppm) As Received	Arsenic (ppm) Washed & Dried In Kitchen	Arsenic (ppm) Cleaned As In Kitchen
14.	Mr. A. P. Morris	0.61		0.11
	'	0.15		0.11
	"			0.04
	"	0.54		0.08
	"	0.23		0.13
15.	Correctional Camp			0.02
	"	0.32		0.01
	"			0.09
	"			0.01
	"	0.25		0.03
16.	Mr. Christensen-Old Town	0.49		0.07
	"	0.34		0.06
	"			0.49
	"	0.11		0.06
17.	5018-54 Street, Yellowknife	0.73		0.77
	"	0.07		0.25

TABLE 34

Arsenic Content in Yellowknife Vegetables,
September, 1976.
(Reference 15)

Location	Vegetable	Total Arsenic (ppm) (natural weight basis)
W.A. Case residence (Yellowknife)	Potatoes	< 0.02
D.H. Egli residence (Yellowknife)	Lettuce	0.21
	Rhubarb	0.04
	Carrots	0.07
	Potatoes	0.10
	Beets	-
	Cabbage	0.29
Con Gardens	Lettuce	0.29
	Rhubarb	0.11
	Carrots	0.11
	Potatoes	0.03
	Beets	-
	Turnips	-
	Cabbage	0.06
Giant Gardens (various locations)	Lettuce	-
	Rhubarb	0.16
	Carrots	0.20
	Potatoes	0.08
	Beets	-
	Turnips	0.11
	Parsley	0.79

Notes: (1) All vegetables were thoroughly cleaned and washed; weighed in natural state; dried at 100°F to remove all moisture; and analysed for arsenic. Analyses are calculated on natural weight.

(2) Analyses conducted at Trail Laboratory (arsine generation method).

TABLE 35

Arsenic Content of Native Grass Samples,
Yellowknife, N.W.T.

Mid-summer Surveys

(Reference 15)

Location	Total Arsenic (ppm)		
	1970	1975	1976
Con Property - 100 yards N. of Robertson headframe	74.0	44.0	32.0
Negus Mine - 100 yards N.E. of old shaft	8.0	22.0	139.0
Con Gardens	8.0	12.0	-
Rat Lake - 25 yards W. of lake	-	6.0	10.0
Yellowknife - D.H. Egli residence	-	10.0	-
Frame Lake - N. of lake	20.0	10.0	8.0
Giant - Edmonton road junction - 75 yards N.	6.0	28.0	16.0
Giant Mine - 1 mile S. of stack	11.0	17.0	21.0

- Notes:
- (1) Samples collected by cutting the grass about one inch above roots.
 - (2) Samples air dried prior to analysis.
 - (3) Analyses are on a dry weight basis.
 - (4) Analyses conducted at Trail laboratory (arsine generation method).

TABLE 36

Arsenic in Fish Livers and Fish Muscles

Falk (11)

1973

Location	Fish Type	Number of Fish Sampled	Average Arsenic Concentration (ppm wet weight)	
			Liver	Muscle
Near mouth of Baker Creek	Sucker	4	--	0.28
	Whitefish	1	0.87	0.22
	Northern Pike	4	0.70	0.31
Back Bay, west of Latham Island.	Whitefish	2	<0.20	<0.20
	Northern Pike	1	0.73	--
Martin Lake (head- water of Baker Creek)	Northern Pike	5	0.96	1.69
	Walleye	4	1.62	0.29
	Whitefish	4	0.53	0.28
Kam Lake	Whitefish	5	0.80	0.28
	Northern Pike	5	2.23	3.22
Yellowknife Bay, sub-island region.	Northern Pike	5	0.32	0.31
	White fish	3	0.24	<0.20

<u>Location</u>	<u>Animal</u>	<u>Arsenic Concentration, ppm</u>
Keg Lake	Snails	79.7
Kam Lake	Snails	48.6
Vee Lake	Snails	4.8
Madeline Lake	Snails	3.1
Near Latham Island	Snails	5.4
Graham Lake	Water Beetles, amphipods	3.5

TABLE 37

Arsenic in Fish Muscle
 Fisheries & Marine Service,
 Yellowknife, N.W.T. (12)
 1975

Location	Fish Type	Number of Fish Sampled	Arsenic Concentration (ppm)
Back Bay, north of Baker Creek.	Northern Pike	6	<0.20
	Whitefish	2	<0.20
Back Bay, south of Baker Creek.	Whitefish	3	<0.20
	Maria	2	<0.20
East Mirage Islands, Yellowknife Bay.	Whitefish	4	<0.20
	Maria	1	<0.20

TABLE 38

Number of Organisms

In Yellowknife Area Lakes and Rivers

Baker CreekAverage summer densities ($\pm 95\%$ confidence limits)

	<u>Above Mine</u>	<u>Below Mine</u>
Phytoplankton	$0.8 \pm 0.1 \times 10^8 \mu\text{m}^3/\text{L}$	0.003×10^8
Epiphyton	$2.5 \pm 0.2 \times 10^8 \mu\text{m}^3/\text{cm}^2$	0.0009×10^8
Rotiferan zooplankton	500 animals/ m^3	0
Crustacean zooplankton	12/ m^3	0
Zoobenthos	5500/ m^2	<100/ m^2

Yellowknife Bay

(a) Transect directly out from Giant Mine into Yellowknife Bay

<u>Distance from wharf</u> <u>(m)</u>	<u>Animals/m^2</u>
At wharf	0.00
200	550
400	200
600	1100
800	3600) *
1000	6600)
1200	4500)
1400	1100
1600	3900
1800	900

* recovery in numbers occurs in an area where the bottom is elevated and therefore protected from tailings.

(b) From Giant Mine into Back Bay

At wharf	0
200	2300
400	1500
600	200
800	250
1000	200
1200	400
1400	200
1600	900
1800	1600
2000	3100
2200	6700

TABLE 39

Arsenic Content of Small Wildlife
Game Management Branch,
Government of the Northwest Territories

April, 1976
(values are in ppm arsenic)
(whole body analysed)
(Reference 13)

Specimen Number	Specimen	Collection Date	Water Content (%)	Arsenic Concentration	Collection Location
1	Willow Ptarmigan	Feb. 18/76	71.38	0.75	on road at Vee Lake
2	"	"	71.07	0.72	"
3	"	"	71.84	0.45	"
4	"	"	71.20	0.59	"
5	Snowshoe Hare	Feb. 19/76	75.53	1.44	Giant Mine dump.
6	Raven	"	69.79	<0.1	Bow-Springer kennels.
7	"	"	69.23	<0.1	"
8	"	"	69.42	<0.1	"
9	"	"	70.07	<0.1	"
10	"	"	67.52	<0.1	"
11	Willow Ptarmigan	Feb. 20/76	71.01	0.34	on road at Vee Lake
12	"	"	71.45	0.35	"
13	"	"	71.01	0.44	"
14	"	"	70.79	0.32	"
15	"	Nov. 12/75	69.31	<0.1	Highway 2; 26 miles west of Yellowknife.

Note: Analyses conducted at Canadian Wildlife Service, Ottawa.

TABLE 40

Arsenic in Human Hair Samples

Yellowknife, N.W.T.
(Reference 14)
1975

Early in 1975, 700 Yellowknife residents volunteered to have their hair sampled and tested for arsenic content.

Number of Persons with Arsenic Concentration

	<u>Under 5 ppm</u>	<u>5-10 ppm</u>	<u>Over 10 ppm</u>	<u>Total</u>
Mine & Mill Workers	61	30	44	135
Other Residents	<u>516</u>	<u>30</u>	<u>19</u>	<u>565</u>
All persons tested	577	60	63	700

Fifty-seven people from the "over 10 ppm" group received a thorough medical examination. Arsenic content of hair samples of this group were:

<u>Arsenic in Hair (ppm)</u>	<u>Number of Persons</u>	<u>Arsenic in Hair mean value, ppm</u>
10	10	7.1
10 - 49	34	21.2
50 - 99	7	66.7
Greater than 100	<u>6</u>	<u>203.0</u>
	57	43.5

TABLE 41

Arsenic in Human Urine Samples

Cominco Con Mine Personnel
Yellowknife, N.W.T.

February-March 1977

(Reference 16)

Group	Number of Employees with Urinary Arsenic Concentration										Total
	< 20	20-30	30-40	40-50	50-60	60-70	70-80	80-90	90-100	> 100	
(1) Under-ground	85	17	8	2	1	0	1	-	-	-	114
(2) Mechanical & Trades	30	2	0	2	-	-	-	-	-	-	34
(3) Office	26	2	0	1	-	-	-	-	-	-	29
(4) Mill Personnel	10	5	0	1	1	1	1	1	-	5	25
(5) Misc. Surface	9	2	-	-	-	-	-	-	-	-	11

Totals	160	28	8	6	2	1	2	1	0	5	213
(%)	(75.1)	(13.2)	(3.8)	(2.8)	(0.9)	(0.5)	(0.9)	(0.5)	(0)	(2.4)	(100)

(N.B. Arsenic analyses are in units of ugm As/litre)

TABLE 41

Arsenic in Human Urine Samples

Cominco Con Mine Personnel
Yellowknife, N.W.T.

February-March 1977

(Reference 16)

Notes:

In February 1977 a voluntary urinary arsenic survey was conducted by Cominco Limited on 213 out of a total of 236 employees at the Con operation in Yellowknife.

Twenty-three employees did not submit specimens either due to personal reasons, sickness, vacation, resignation or attending schools out of town.

- (a) Can Test Limited (Vancouver) supplied specimen bottles and performed the analyses.
- (b) Specimen bottles were issued February 21-22, 1977, and collected over a three-week period.
- (c) Two of the five employees reporting levels greater than 100 ug/m As/litre have lived in Yellowknife and worked at Con Mine less than two years.
- (d) For reference purposes, normal urinary arsenic values for persons not exposed to arsenic, range from 10 to 300 ug/m As per litre.

REFERENCES

1. Toft, P., Bogoroch, R., Sandi, E., Sundaram, A., and Windish, J.P.,
"Report of the Task Force on Arsenic in Ambient Air",
Environmental Health Directorate, Department of
National Health and Welfare, Ottawa, July 1975.
2. Capowski, R., "Yellowknife Arsenic Stack Survey", Air
Pollution Control Section, E.P.S., Northwest Region,
Edmonton, (June 1977).
3. Hazra, A.K., and Prokopuk, R., "Report on the Air Quality
Levels in the Yellowknife Area", 1975, Air Pollution
Control Section, E.P.S., Northwest Region, Edmonton,
(June 1977).
4. O'Toole, J.J., Clark, R.G., Malaby, K.L., and Trauger, D.L.,
"Environmental Trace Element Survey at a Heavy Metals
Refining Site", 1970.
5. de Villiers, A.J., and Baker, P.M., "An Investigation of the
Health Status of Inhabitants of Yellowknife, N.W.T.",
Department of National Health and Welfare, Ottawa, 1969.
6. Hazra, A.K., and Hardin, M.J., "Report on the Quality of Snow in
the Yellowknife Area", 1975, Air Pollution Control Section,
E.P.S., Northwest Region, Edmonton, June 1977.
7. Martin, R.W., "Arsenic Concentrations in Soil and Road Dust
Samples taken from Areas Utilized by the Inhabitants of
Yellowknife", E.P.S., Northwest Region, Yellowknife, June 1977.
8. Temple, P.J., Linzen, S.N., and Chai, B.L., "Contamination of
Vegetation and Soil by Arsenic Emissions from Secondary
Lead Smelters" unidentified

9. Brown, R.L., Supervisor, Waste Control, Cominco Ltd., Trail, B.C.,
internal correspondence, Yellowknife Arsenic Survey,
November 12, 1975.
10. Plambeck, J.A. and Smith, R.A., "Yellowknife Environmental Survey -
Soils and Vegetation", Report prepared for E.P.S., Environment
Canada, (April, 1976).
11. Falk, M.R., Miller, M.D., and Kostuik, S.J.M., "Biological Effects
of Mining Wastes in the NWT", Fisheries and Marine Service,
Environment Canada, Resource Management Branch, Report
number CEN/T-73-10, (October, 1973).
12. Glazier, G.T., District Manager, Fisheries and Marine Service,
Yellowknife, to W.J. Bryant, District Manager, E.P.S.,
Yellowknife, May, 1975: Arsenic Analyses in Fish Samples.
13. Canadian Wildlife Service, Ottawa, correspondence to Mr. D.
Billing, Environmental Protection Officer, Government of
the Northwest Territories, Yellowknife, April, 1976:
Arsenic Determinations in Small Wildlife.
14. Schaefer, O., "Effects of Arsenic on Human Health in the
Yellowknife Area", Department of National Health and
Welfare, July, 1975.
15. Cominco Limited, Con Operations, "Arsenic Survey 1976",
Yellowknife, N.W.T., May, 1977.
16. Cominco Limited, Con Operations, "1977 Urinary Arsenic Study".
Yellowknife, N.W.T., May, 1977.

